arXiv:cond-mat/0610380v1 [cond-mat.other] 13 Oct 2006

Ultracold heteronuclear molecules and ferroelectric superfluids

M. Iskin and C. A. R. Sá de Melo
School of Physics, Georgia Institute of Technology, Atlanta, Georgia 30332, USA

(Dated: February 6, 2008)

We analyze the possibility of a ferroelectric transition in heteronuclear molecules consisting of Bose-Bose, Bose-Fermi or Fermi-Fermi atom pairs. This transition is characterized by the appearance of a spontaneous electric polarization below a critical temperature. We discuss the existence of a ferroelectric Fermi liquid phase for Fermi molecules and the existence of a ferroelectric superfluid phase for Bose molecules characterized by the coexistence of ferroelectric and superfluid orders. Lastly, we propose an experiment to detect ferroelectric correlations through the observation of coherent dipole radiation pulses during time of flight.

PACS numbers: 03.75.Kk, 03.75.Ss, 32.10.Dk, 33.15.-e

Arguably one of the next frontiers in ultracold atomic and molecular physics is the study of two-species atomic mixtures [1, 2, 3, 4, 5, 6], and ultracold heteronuclear molecules composed of two-species alkali atoms such as KRb [7, 8, 9], RbCs [10] and NaCs [11]. This frontier may be advanced through the use of Feshbach resonances which have already been observed in mixtures of two-species alkali atoms [12, 13, 14], and may also become a crucial tool for tuning physical properties of heteronuclear systems.

Ultracold heteronuclear molecules made of Bose-Bose, Bose-Fermi or Fermi-Fermi atom pairs offer many new opportunities compared to standard (Bose or Fermi) atomic systems because of their additional degrees of freedom. For instance, when heteronuclear diatomic molecules are formed from neutral atoms, electric charge is transferred from one atom to the other leading to an electric dipole moment $|\mathbf{p}| = Q(d)d$, where d is the separation and $Q(d)$ is the effective charge transfer between constituent atoms [15]. These electric dipoles have equal magnitudes but random orientations at high temperatures leading to a vanishing average electric polarization. However, at low temperatures, the dipoles may all point to a particular direction producing a finite average electric polarization density $\langle \mathbf{P} \rangle$, characteristic of a ferroelectric state. In addtion, when ultracold heteronuclear molecules form a Bose-Einstein condensate (BEC) a ferroelectric superfluid state proposed in this manuscript may be accessible experimentally.

Our main results are as follows. When heteronuclear molecules are composite fermions (Bose-Fermi pairs), a phase transition occurs separating a paraelectric Fermi liquid (PFL) from a ferroelectric Fermi liquid (FFL), as shown in Fig. 1(a). These phases do not exist in standard condensed matter systems, since all experimentally known ferroelectrics are very good insulators [16], and are not Fermi liquids. Furthermore, when heteronuclear molecules are composite bosons (Bose-Bose or Fermi-Fermi pairs), a ferroelectric transition with critical temperature T_{FE} may occur either above or below the BEC temperature T_{BEC}. When the molecular dipole moment and/or density are sufficiently large then $T_{FE} > T_{BEC}$ leading to a paraelectric Bose liquid (PBL) for $T > T_{FE}$, a ferroelectric Bose liquid (FBL) for $T_{FE} > T > T_{BEC}$, and to a ferroelectric BEC (FBEC) for $T < T_{BEC}$, as shown in Fig. 1(b). When the molecular dipole moment and/or density are sufficiently low then $T_{FE} < T_{BEC}$ leading to a paraelectric Bose liquid (PBL) for $T > T_{BEC}$, a paraelectric BEC for $T_{BEC} > T > T_{FE}$, and to a ferroelectric BEC (FBEC) for $T < T_{FE}$, as shown in Fig. 1(c). The FBEC phase corresponds to a ferroelectric superfluid.



FIG. 1: Schematic phase diagram for polar a) Fermi, and b), c) Bose molecules, where $\langle \mathbf{P} \rangle$ and $\langle \phi \rangle$ are the corresponding order parameters for ferroelectric order at T_{FE}, and for BEC at T_{BEC}, respectively.

Hamiltonian: To describe all these phases of ultracold heteronuclear molecules, we use the Hamiltonian density

$$\begin{aligned} H(\mathbf{r}) &= \sum_{\sigma} \psi^*_{\sigma}(\mathbf{r}) \left[-\frac{\nabla^2}{2M} - \mu + V(\mathbf{r}) \right] \psi_{\sigma}(\mathbf{r}) \\ &+ \frac{1}{2} \sum_{\sigma,\sigma'} \int d\mathbf{r}' n_{\sigma'}(\mathbf{r}) U_d(\mathbf{r},\mathbf{r}') n_{\sigma}(\mathbf{r}) \\ &- \frac{1}{2} \sum_{\alpha,\beta} \int d\mathbf{r}' \mathbf{P}_{\alpha}(\mathbf{r}') J_{\alpha,\beta}(\mathbf{r},\mathbf{r}') \mathbf{P}_{\beta}(\mathbf{r}), \qquad (1) \end{aligned}$$

describing a weakly interacting gas of dilute polar molecules, where $\psi^*_\sigma(\mathbf{r})$ creates a Bose (Fermi) molecule at position $\mathbf{r}$ with pseudo-spin-σ, and $n_\sigma(\mathbf{r}) = \psi^*_\sigma(\mathbf{r})\psi_\sigma(\mathbf{r})$ is the density operator. The first term in Eq. (1) is the kinetic energy, where μ is the chemical potential, and $V(\mathbf{r})$ is the trapping potential. The second term is the density-density interaction between Bose (Fermi) molecules, and for a contact interaction is given by $U_d(\mathbf{r},\mathbf{r}') = U\delta(\mathbf{\Delta r})$, where $U = 4\pi a/M$ is the amplitude of the interaction and $\delta(\mathbf{\Delta r})$ is the delta function with $\mathbf{\Delta r} = \mathbf{r} - \mathbf{r}'$. Here, a is the scattering length of the corresponding Bose (Fermi) molecules, and M is the molecular mass. The third term is the electric dipole-dipole interaction between molecular dipoles at positions $\mathbf{r}$ and $\mathbf{r}'$, where $\mathbf{P}(\mathbf{r}) = \mathbf{p}(\mathbf{r})\sum_\sigma n_\sigma(\mathbf{r})$ is the polarization operator, and $\mathbf{p}(\mathbf{r})$ is the molecular dipole moment at position $\mathbf{r}$. The electric dipole-dipole interaction is given by $J(\mathbf{r},\mathbf{r}') = \sum_{\alpha,\beta}\mathbf{p}_\alpha(\mathbf{r})J_{\alpha,\beta}(\mathbf{r},\mathbf{r}')\mathbf{p}_\beta(\mathbf{r}')$, where

$$J_{\alpha,\beta}(\mathbf{r},\mathbf{r}') = \frac{1}{4\pi\varepsilon_0}\frac{3[\mathbf{\Delta r}_\alpha \mathbf{\Delta r}_\beta]/|\mathbf{\Delta r}|^2 - \delta_{\alpha,\beta}}{|\mathbf{\Delta r}|^3}. \quad (2)$$

Here, $\delta_{\alpha,\beta}$ is the Kronecker-delta and $\{\alpha,\beta\} = \{1,2,3\}$ label the vector components.

Electric polarization: Within the Hartree-Fock approximation, the Hamiltonian density reduces to

$$\begin{aligned} H_0(\mathbf{r}) &= -\frac{\nabla^2}{2M} - \widetilde{\mu} + V(\mathbf{r}) + Un_0(\mathbf{r}) \\ &- \sum_{\alpha,\beta}\int d\mathbf{r}'\mathbf{p}_\alpha(\mathbf{r})J_{\alpha,\beta}(\mathbf{r},\mathbf{r}')\mathbf{P}_{0,\beta}(\mathbf{r}'), \end{aligned} \quad (3)$$

where the electric polarization density $\mathbf{P}_0(\mathbf{r}) = \mathbf{p}(\mathbf{r})n_0(\mathbf{r})$, with $n_0(\mathbf{r}) = \sum_\sigma n_{0,\sigma}(\mathbf{r})$ being the local density of Bose (Fermi) molecules at $\mathbf{r}$. The spatially averaged polarization $\langle\mathbf{P}\rangle = \int d\mathbf{r}\mathbf{P}_0(\mathbf{r})/V_c$ can be rewritten as

$$\langle\mathbf{P}\rangle = \frac{1}{V_c}\int d\mathbf{r}\mathbf{p}(\mathbf{r})\sum_{i,\sigma}|\phi_i(\mathbf{r})|^2 f_\eta(\epsilon_i), \quad (4)$$

where V_c is the sample volume, $\phi_i(\mathbf{r})$ and ϵ_i are eigenfunctions and eigenvalues of the Hamiltonian $H_0(\mathbf{r})$, and $f_\eta(\epsilon_i) = 1/[e^{\beta\epsilon_i} - \eta]$ is the Bose (Fermi) distribution for Bose (Fermi) molecules when $\eta = 1$ (-1). The solution of Eq. (4) is non-trivial, however analytical insight can be gained for homogenous systems where $V(\mathbf{r}) = 0$ and $\mathbf{P}_0(\mathbf{r}) = \mathbf{P}_0$ is independent of $\mathbf{r}$. We discuss the homogeneous case first and then analyze the case of a harmonic trap.

Ferroelectric critical temperature: In the ferroelectric state, all molecular dipoles are pointing along the same direction $\widehat{\mathbf{m}}$ such that $\mathbf{p}(\mathbf{r}) = \mathbf{p} = |\mathbf{p}|\widehat{\mathbf{m}}$ and $\mathbf{P}_0 = |\mathbf{P}_0|\widehat{\mathbf{m}}$. The critical temperature T_{FE} for the ferroelectric transition is found from the slope of Eq. (4) with respect to $\mathbf{P}_0$ evaluated at $\mathbf{P}_0 = 0$, leading to

$$1 + \widetilde{J}(\mathbf{q}\to\mathbf{0})\left(\frac{\partial N}{\partial\mu}\right)_T = 0, \quad (5)$$

from which T_{FE} can be calculated. Here, $\widetilde{J}(\mathbf{q})$ with $\mathbf{q} = \mathbf{k} - \mathbf{k}'$ is the Fourier transform of $J(\mathbf{\Delta r})$, and N is the number of Bose or Fermi molecules. Introducing the Kac parameter γ as a cutoff for short distances ($|\mathbf{\Delta r}| < 1/\gamma$) leads to $\widetilde{J}(\mathbf{q}) = \sum_{\alpha,\beta}\mathbf{p}_\alpha\widetilde{J}_{\alpha,\beta}(\mathbf{k},\mathbf{k}')\mathbf{p}_\beta$, where

$$\widetilde{J}_{\alpha,\beta}(\mathbf{k},\mathbf{k}') = \frac{S(\widetilde{q})}{\varepsilon_0}\left(\frac{3\mathbf{q}_\alpha\mathbf{q}_\beta}{|\mathbf{q}|^2} - \delta_{\alpha,\beta}\right). \quad (6)$$

Here, $S(\widetilde{q}) = \sin\widetilde{q}/\widetilde{q}^3 - \cos\widetilde{q}/\widetilde{q}^2$ where $\widetilde{q} = |\mathbf{q}|/\gamma$. Thus, we obtain an implicit relation for the T_{FE}

$$1 - \frac{2|\mathbf{p}|^2}{3\varepsilon_0}n^2\kappa(T_{FE}) = 0, \quad (7)$$

in terms of the molecular density $n = N/V_c$ and the isothermal compressibility $\kappa(T) = (1/n^2)(\partial n/\partial\mu)_T$.

The ferroelectric instability is accompanied by a divergence of the dielectric function in the long-wavelength and low-frequency limit. Using linear response theory, the dielectric function $\varepsilon(\mathbf{q}, iw_n)$ can be related to the density-density correlation function $C(\mathbf{q},\tau) = \langle T_\tau n(\mathbf{q},\tau)n(-\mathbf{q},0)\rangle$ and to $\widetilde{J}(\mathbf{q})$ via

$$\frac{1}{\varepsilon(\mathbf{q}, iw_n)} = 1 - \frac{\widetilde{J}(\mathbf{q})}{V_c}\int_0^{1/T} d\tau e^{iw_n\tau}C(\mathbf{q},\tau), \quad (8)$$

where $n(\mathbf{q},\tau)$ is the density operator. In the long-wavelength and low-frequency limit, $C(\mathbf{q},\tau)$ is directly related to $\kappa(T)$ via the compressibility sum rule [17]. Therefore, a divergent dielectric function occurs when $\varepsilon(\mathbf{q}\to 0, iw_n \to 0) = 0$ leading to $1 + \widetilde{J}(\mathbf{q}\to 0)(\partial N/\partial\mu)_T = 0$, which is identical to Eq. (5). This relation can be applied to both Bose and Fermi systems. Next, we discuss T_{FE} for a weakly interacting ($a \ll \lambda$) and dilute ($na^3 \ll 1$) gas of Bose and Fermi molecules, where a is the scattering and λ is the thermal length.

Fermi molecules: As a first application of Eq. (7), we analyze T_{FE} for a weakly interacting gas of Fermi molecules at any T. In this case, the molecular density is given by $n = 2F_{3/2}(z)[1 - 2F_{1/2}(z)a/\lambda_F]/\lambda_F^3$ where $\lambda_F = [2\pi/(MT)]^{1/2}$ is the thermal length, and leading to $\kappa(T) \approx [F_{1/2}(z)/F_{3/2}(z) - 2F_{-1/2}(z)a/\lambda_F]/(nT)$ where $0 \le z = \exp(\beta\mu) \le \infty$ is the fugacity, and $F_\nu(z) = [1/\Gamma(\nu)]\int_0^\infty x^{\nu-1}dx/[z^{-1}e^x + 1]$ is the Fermi integral. Here, $\Gamma(\nu)$ is the Gamma function. Thus, we obtain

$$T_{FE} = \frac{2|\mathbf{p}|^2}{3\varepsilon_0}\left[\frac{F_{1/2}(z_c)}{F_{3/2}(z_c)} - \frac{2a}{\lambda_F}F_{-1/2}(z_c)\right]n, \quad (9)$$

where $z_c = \exp(\mu/T_{FE})$. Notice that, in the classical ($z_c \ll 1$) limit, Eq. (9) reduces to $T_{FE} = 2|\mathbf{p}|^2(1 - 2z_c a/\lambda_F)n/(3\varepsilon_0)$, which shows that $T_{FE} \propto n$ for a gas of classical electric dipoles. The T_{FE} for an ideal (non-interacting) gas of Fermi molecules can be obtained by setting $a = 0$. For an ideal gas, when T_{FE}

is much smaller than the Fermi energy ϵ_F, we obtain $T_{FE} \approx (2\sqrt{3}/\pi)\epsilon_F[|\mathbf{p}|^2 n/(\varepsilon_0\epsilon_F) - 1]^{1/2}$, which is valid for $|\mathbf{p}|^2 n/(\varepsilon_0\epsilon_F) > 1$ and $[|\mathbf{p}|^2 n/(\varepsilon_0\epsilon_F) - 1]^{1/2} \ll 1$. For $T > T_{FE}$ a PFL phase exists and for $T < T_{FE}$ a FFL phase appears as shown in Fig. 1(a).

Bose molecules for $T \geq T_{BEC}$: As a second application of Eq. (7), we analyze T_{FE} for a weakly interacting gas of Bose molecules when $T \geq T_{BEC}$. In this case, the molecular density is given by $n = B_{3/2}(z)[1 - 4B_{1/2}(z)a/\lambda_B]/\lambda_B^3$ where $\lambda_B = [1/(2\pi MT)]^{1/2}$ is the thermal length, and leading to $\kappa(T) \approx [B_{1/2}(z)/B_{3/2}(z) - 4B_{-1/2}(z)a/\lambda_B]/(nT)$ where $0 \leq z = \exp(\beta\mu) \leq 1$ is the fugacity, and $B_\nu(z) = [1/\Gamma(\nu)]\int_0^\infty x^{\nu-1}dx/[z^{-1}e^x - 1]$ is the Bose integral. Thus, we obtain

$$T_{FE} = \frac{2|\mathbf{p}|^2}{3\varepsilon_0}\left[\frac{B_{1/2}(z_c)}{B_{3/2}(z_c)} - \frac{4a}{\lambda_B}B_{-1/2}(z_c)\right]n. \qquad (10)$$

Notice that, in the classical ($z_c \ll 1$) limit, Eq. (10) reduces to $T_{FE} = 2|\mathbf{p}|^2(1 - 4z_c a/\lambda_B)n/(3\varepsilon_0)$, which again leads to $T_{FE} \propto n$ for a gas of classical electric dipoles. The T_{FE} for an ideal (non-interacting) gas of Bose molecules can be obtained by setting $a = 0$. Notice that, the second terms in Eqs. (9) and (10) are different by a factor of 2 due to the degeneracy of pseudo-spin-1/2 fermions in contrast to pseudo-spin-0 bosons.

Bose molecules for $T \leq T_{BEC}$: As a third application of Eq. (7), we analyze T_{FE} for an ideal (non-interacting) gas of Bose molecules when $T \leq T_{BEC}$. In this case, the molecular density is given by $n = B_{3/2}(z)/\lambda_B^3 + n_s$ where $n_s = z/[V_c(1-z)] = \alpha(T)n$ is the density of bosons in the condensed (zero-energy) state. Here, $\alpha(T) = 1 - (T/T_{BEC})^{3/2}$ where $T_{BEC} = 2\pi[n/\zeta(3/2)]^{2/3}/M$ is the critical BEC temperature for non-interacting dilute bosons, and $\zeta(x)$ is the Zeta function. In this case, $\kappa(T) = [B_{1/2}(z)/\lambda_B^3 + V_c n_s^2/z]/(n^2 T)$, which diverges in the thermodynamic limit [18] when $\{N, V_c\} \to \infty$ but $n = N/V_c$ is a constant. Thus, we obtain

$$T_{FE} = \frac{2|\mathbf{p}|^2}{3\varepsilon_0}\left[\frac{B_{1/2}(z_c)}{B_{3/2}(z_c)}(n - n_s) + \frac{V_c n_s^2}{z_c}\right], \qquad (11)$$

which is always smaller than T_{BEC}, and reduces to the non-interacting limit of Eq. (10) in the absence of BEC.

The two cases of Bose molecules allows the construction of the phase diagrams indicated in Fig. 1(b) and 1(c), respectively, where the PBL, FBL, PBEC and FBEC are identified depending on the existence of a spontaneous average electric polarization $\langle\mathbf{P}\rangle$ and/or of a BEC fraction $\langle\phi\rangle$. The ferroelectric superfluid phases proposed here may be experimentally observed with currently available cooling techniques only when T_{FE} is large enough. This requirement imposes a condition on the size of the electric dipole moments of the molecules, and it is discussed next for the Fermi-Fermi Bose molecules.

Fermi-Fermi Bose molecules: To set the scale, we consider the specific example of Li-K molecules consisting of ^{6}Li and ^{40}K atoms in their ground state, where $|\mathbf{p}| = 3.6$ Debye [15]. We also choose an equal population mixture of ^{6}Li and ^{40}K atoms with parameters $N = 10^5$ and $V = 10^{-7}$ cm^3 leading to $T_{BEC} \approx 0.099\epsilon_F$ and $T_{FE} \approx 21T_{BEC}$. However, for a molecule with $|\mathbf{p}| = 1.0$ Debye, $T_{FE} = 1.6T_{BEC}$. Here, $\epsilon_F = k_F^2/(2m_r)$ is the Fermi energy, where m_r is half of the reduced mass of Li and K atoms, and k_F is the Fermi momentum with $n = k_F^3/(6\pi^2)$.



FIG. 2: Schematic phase diagram of temperature T/T_{BEC} versus scattering parameter $1/(k_F a)$ for an equal mixture of two-species Fermi-Fermi mixtures.

For the two-species Fermi-Fermi mixtures (such as ^{6}Li and ^{40}K), we propose a schematic phase diagram of temperature T versus scattering parameter $1/(k_F a)$ as shown in Fig. 2. In the diagram, T_0 is the pair formation scale and Cooper pairs break above this temperature. In addition, a two-body bound state does not exist for negative a. Thus, a polar bosonic molecule with dipole moment $|\mathbf{p}| \neq 0$ may form when both $T < T_0$ and $a > 0$ criteria are satisfied. This also implies that a finite polarization $\mathbf{P}_0$ may exist below $T < T_{FE}$ in the phase space where $T < T_0$ and $a > 0$ leading to a possible ferroelectric phase. Depending on the relative values of T_{FE} and T_{BEC}, we sketched two cases: (I) $T_{FE} > T_{BEC}$ and (II) $T_{FE} < T_{BEC}$ which are shown as dotted lines. Notice that as $1/(k_F a)$ decreases, the size of the molecules increases and $\mathbf{p}$ gets smaller, thus producing a reduced T_{FE}. Therefore, an equal mixture of ^{6}Li and ^{40}K correspond to case I in Fig. 2. However, other mixtures with smaller dipole moments and/or densities may correspond to case II. Next, we discuss trapped molecules.

Heteronuclear molecules in a trap: The trapping potential of polar molecules is given by $V(\mathbf{r}) = -\alpha|\mathbf{E}(\mathbf{r})|^2 - \mathbf{p}\cdot\mathbf{E}(\mathbf{r})$, which are due to the coupling of the laser electric field $\mathbf{E}(\mathbf{r})$ with the induced and permanent dipole moments $\mathbf{p}_{ind} = \alpha\mathbf{E}(\mathbf{r})$ and $\mathbf{p}_{per} = \mathbf{p}$, respectively. We assume that $\mathbf{E}(\mathbf{r}) = E_0\hat{\mathbf{x}}\exp[-\rho^2/(2w_\rho^2) - z^2/(2w_z^2)]$ has a gaussian profile controlled by the widths w_ρ and w_z, where $\rho^2 = x^2 + y^2$. Thus, the trapping potential can be approximated by $V(\mathbf{r}) = V_0 + M\Omega_\rho^2\rho^2/2 + M\Omega_z^2 z^2/2$, where $V_0 = -\alpha E_0^2 - |\mathbf{p}|E_0 \cos\theta$, and $\Omega_i = (2\alpha E_0 + |\mathbf{p}|E_0 \cos\theta)/(M w_i^2)$ are the characteristic frequencies of the harmonic trap along $i = \rho, z$ directions. Here, θ is the angle between $\mathbf{p}$ and $\mathbf{E}(\mathbf{r})$. When $w_z \gg w_\rho$ ($w_z \ll w_\rho$) the trap is cigar (disc) shaped.

In the ground state of the system, the Thomas-Fermi approximation can be applied leading to

$$U n_0(\mathbf{r}) = \widetilde{\mu} - V(\mathbf{r}) + \int d\mathbf{r}' J(\mathbf{r}, \mathbf{r}') n_0(\mathbf{r}'), \qquad (12)$$

and the electric polarization density is given by $\mathbf{P}(\mathbf{r}) = \mathbf{p} n_0(\mathbf{r})$. The integral equation can be solved analytically when $\beta = [|\mathbf{p}|^2/(4\pi\varepsilon_0)]/U \ll 1$, corresponding to a small ratio between the characteristic electric dipolar energy $|\mathbf{p}|^2/(4\pi\varepsilon_0 V_c)$ and the characteristic contact interaction energy U/V_c. To zeroth order in β, the electric polarization is $\mathbf{P}(\mathbf{r}) = \mathbf{P}_{TF}(\mathbf{r}) = \mathbf{p} n_{TF}(\mathbf{r})$, where

$$n_{TF}(\mathbf{r}) = n_{max} \left[1 - \rho^2/\rho_c^2 - z^2/z_c^2\right]. \qquad (13)$$

Here, $n_{max} = (\widetilde{\mu} - V_0)/U$, $\rho_c^2 = (\widetilde{\mu} - V_0)/(M\Omega_\rho^2)$ and $z_c^2 = (\widetilde{\mu} - V_0)/(M\Omega_z^2)$. The correction to first order in β for the cigar shaped trap is

$$\delta\mathbf{P}(\mathbf{r}) = \mathbf{p}\beta \frac{4\pi}{18} \bar{n} \frac{\rho_c^2}{z_c^2} \left[c_1 - c_2(\rho^2/\rho_c^2 - z^2/z_c^2)\right], \qquad (14)$$

where c_1 and c_2 are numerical coefficients, and $\bar{n} = N/V_c$ is the average molecular density. Since $\mathbf{E}(\mathbf{r})$ is much stronger than the local electric fields produced by electric dipolar interactions, the net polarization $\mathbf{P}(\mathbf{r}) = \mathbf{P}_{TF}(\mathbf{r}) + \delta\mathbf{P}(\mathbf{r})$ points along $\mathbf{E}(\mathbf{r})$.



FIG. 3: Direction of the polarization vector during time of flight for a ferroelectric state, when the external electric field is a) perpendicular, and b) parallel to the easy axis of polarization. The clouds in each frame are not drawn to scale.

To distinguish between the fully polarized paraelectric superfluid and the true ferroelectric superfluid, we propose two experiments, as shown schematically in Fig. 3, where $\mathbf{E}(\mathbf{r})$ is turned off between times I and II. In case (a), the external electric field and the easy axis for polarization are perpendicular. For the ferroelectric state, the polarization flips between times I and II (within nanoseconds to microseconds) towards the axial direction which corresponds to the easy axis for the electric polarization. This flip produces a short pulse of coherent dipolar radiation proportional to N^2. However, for the fully polarized paraelectric state the polarization relaxes and reduces to zero within microseconds, producing a longer pulse of incoherent dipolar radiation proportional to N. A second coherent pulse may occur in time of flight for the ferroelectric state as the anisotropy inversion in the cloud expansion takes place between times III and IV, and causes a second flip of the electric polarization towards the new easy axis. In case (b), the external electric field and the easy axis for polarization coincide. For the ferroelectric state, the polarization remains along the easy axis between times I and II, and between times III and IV when the anisotropy inversion occurs, the polarization flips towards the new easy axis with emission of coherent dipolar radiation (proportional to N^2). In this case only one pulse (within nanoseconds to microseconds) should be observed after a few miliseconds of expansion. For the paraelectric state similar results as in case (a) apply.

In conclusion, we analyzed the possibility of a ferroelectric transition in Bose-Bose, Bose-Fermi or Fermi-Fermi heteronuclear molecules. This transition is characterized by the appearance of a spontanous electric polarization below a critical temperature where the dielectric function diverges. We obtained the order parameter equation, evaluated the transition temperature and the electric polarization for ultracold heteronuclear (Bose or Fermi) molecules. We discussed the existence of a ferroelectric Fermi liquid phase for polar Fermi molecules, and the existence of a ferroelectric superfluid phase for polar Bose molecules characterized by the coexistence of ferroelectric and superfluid orders. We also proposed an experiment to detect ferroelectric correlations via the observation of coherent dipole radiation pulses during time of flight. We thank NSF (DMR-0304380) for support.

[1] A. G. Truscott et al., Science **291**, 2570 (2001).
[2] F. Schreck et al., Phys. Rev. Lett. **87**, 080403 (2001).
[3] Z. Hadzibabic et al., Phys. Rev. Lett. **88**, 160401 (2002).
[4] G. Roati et al., Phys. Rev. Lett. **89**, 150403 (2002).
[5] G. Modugno et al., Phys. Rev. Lett. **89**, 190404 (2002).
[6] C. Silber et al., Phys. Rev. Lett. **95**, 170408 (2005).
[7] M. W. Mancini et al., Phys. Rev. Lett. **92**, 13203 (2004).
[8] D. Wang et al., Phys. Rev. Lett. **93**, 243005 (2004).
[9] C. Ospelkaus et al., Phys. Rev. Lett **97**, 120402 (2006).
[10] A. J. Kerman et al., Phys. Rev. Lett. **92**, 153001 (2004).
[11] C. Haimberger et al., Phys. Rev. A **70**, 021402(R) (2004).
[12] C. A. Stan et al., Phy. Rev. Lett. **93**, 143001 (2004).
[13] S. Inouye et al., Phys. Rev. Lett. **93**, 183201 (2004).
[14] F. Ferlaino et. al., Phys. Rev. A **73**, 040702(R) (2006).
[15] M. Aymar and O. Dulieu, J. Chem. Phys. **122**, 204302 (2005).
[16] F. Jona, and G. Shirane, Ferroelectric Crystals (Dover, New York, 1993).
[17] D. Pines and P. Nozieres, *The Theory of Quantum Liquids*, Benjamin, (1966).
[18] R. K. Pathria, *Statistical Mechanics*, Butterworth-Heinemann, (1996).